

Research Article

# Evaluating Cross Sections of Gallium Isotopes Production Using proton and deuteron Irradiation

Raghad S. Mohammed

Department of physics, College of Science, Mustansiriyah University, IRAQ.

\*Email: raghad\_almaliki@yahoo.com

## Article Info

Received  
19/10/2016

Accepted  
17/04/2017

## Abstract

In the present work, the production of the cross sections of three Gallium isotopes:  $^{66}\text{Ga}_{35}$  ( $t_{1/2} = 9.4\text{h}$ ,  $\beta^+ = 4.2\text{MeV}$ ),  $^{67}\text{Ga}_{36}$  ( $t_{1/2} = 3.2617\text{d}$ ,  $\text{EC} = 100\%$ ) and  $^{68}\text{Ga}_{37}$  ( $t_{1/2} = 68\text{min}$ ,  $\beta^+ = 89\%$ ) have been discussed. The Gallium isotopes have important applications in nuclear medicine, particularly in Positron Emission Tomography (PET), Single Photon Emission Tomography (SPET) imaging technique and used in tumors diagnosing. The production of irradiant  $^{66}\text{Ga}$ ,  $^{67}\text{Ga}$  and  $^{68}\text{Ga}$  is made by irradiation of an enriched Zinc target using proton and deuteron charged particles. Utilizing high cyclotron yield and low radionuclide impurities, the optimum cyclotron energy range has been chosen for the production of Gallium isotopes. The cross sections of (p,xn), (p, $\gamma$ ) and (d,xn) reactions for the production of Gallium isotopes have been evaluated depending upon the empirical data taken from EXFOR library, which is belonging to the International Atomic Energy Agency (IAEA). Also the yield for each reaction has been evaluated.

**Keywords:** cross sections, Gallium Isotopes, optimized energy range

## الخلاصة

في العمل الحالي، تم مناقشة انتاج المقاطع العرضية لثلاث نظائر لعنصر الكاليوم:  $^{66}\text{Ga}$  ( $t_{1/2} = 9.4\text{h}$ ,  $\beta^+ = 4.2\text{MeV}$ ),  $^{67}\text{Ga}$  ( $t_{1/2} = 3.2617\text{d}$ ,  $\text{EC} = 100\%$ ) و  $^{68}\text{Ga}$  ( $t_{1/2} = 68\text{min}$ ,  $\beta^+ = 89\%$ ). واستخدامها في تقنية التصوير المقطعي بالانبعاث البوزيتروني (PET) والتصوير المقطعي بأنبيعات فوتون منفرد (SPET)، واستخدامها في تشخيص الاورام. يتم انتاج نظائر الكاليوم المشعة بتشعيع هدف الزنك المخصب باستخدام الجسيمات المشحونة من البروتونات والديترونات باستخدام أعلى انتاجية للمعجل واقل نسبة شوائب للنظائر المشعة، تم تحديد افضل مدى طاقة للمعجل لانتاج نظائر الكاليوم. تم تقييم المقاطع العرضية لتفاعلات (p, $\gamma$ )، (p,xn) و (d,xn) لانتاج نظائر الكاليوم المشع اعتمادا على القيم العملية المتوفرة في مكتبة (EXFOR) التابعة لوكالة الطاقة الذرية الدولية (IAEA). كذلك تم حساب الانتاجية لكل تفاعل.

## Introduction

The radioisotopes have a significant importance in the field of nuclear medicine, specifically in diagnostic imaging including Single Photon Emission Computed Tomography (SPECT) and Positron Emission Tomography (PET), and in the field of nuclear medical therapy. These radioisotopes are produced using reactors through nuclear reactions or accelerators through bombardment reactions using charged particle such as protons, deuterons and alpha particles [1]. The three radioisotopes of Gallium ( $^{66}\text{Ga}$ ,  $^{67}\text{Ga}$  and  $^{68}\text{Ga}$ ) have a great importance in nuclear medicine, especially in PET scan and in radiotherapy

by attaching them with monoclonal antibodies to detect tumors locations, and in investigation of different diseases [2]. Gallium-66 ( $t_{1/2} = 9.49\text{h}$ ,  $E_{\gamma} = 833.5, 1039.3\text{keV}$ ,  $\beta^+$ : 56.5%,  $E_{\text{max}\beta^+} = 4.153\text{MeV}$ ;  $\text{EC}: 43.5\%$ ), has an intermediate half-life make it suitable for PET imaging of bioprocesses with intermediate to slow target tissue uptake [3]. The Gallium-67 ( $t_{1/2} = 3.2617\text{d}$ ;  $\text{EC} = 100\%$ ), is widely used in medical applications due to its ability to emit several Auger electrons with energies (7–8) keV. These electrons may reach the investigated cell nucleus from the cell surface or from cytoplasm; hence it is effective in single cell killing. Gallium-67 emits rays



with energies ranged from 91 to 394keV, which is suitable to be detected by gamma cameras [4]. for single photon imaging in PET scan [5]. Ga-68 ( $t_{1/2} = 68\text{min}$ ,  $\beta^+$ : 89.1%; EC: 10.9%) has distinctive characteristics that made it used in PET scan imaging and in tumor diagnosing[6]. The aim of the present work is to evaluate the experimental data of nuclear reactions induced by proton and deuteron particles on enriched Zn target and selected optimization energy range for the production of  $^{66}\text{Ga}$ ,  $^{67}\text{Ga}$  and  $^{68}\text{Ga}$  with small and/or medium-sized cyclotrons less than 30MeV and improve efforts in this field of production.

**Materials and Methods**

**Cross Sections Calculations**

The experimental data play an important role in the evaluation of nuclear reaction cross sections. Table (1), shows the experimental data that have been published in (EXFOR) library, which belongs to the International Atomic Energy Agency (IAEA) for proton and deuteron induced reactions specific for Zn target [7]. The empirical data that obtained by different authors listed in table (1) are not identical. In the present work the evaluations was made by including a careful analysis for these data by recalculating the energy in steps of the interval (0.01MeV) and the calculated cross sections were plotted using Matlab-8 programming language as shown in Figures (1→5).

**Yield of Calculated Products**

The yield of a nuclear reaction can be defined as the ratio of the number of the nucleus formed in a nuclear reaction to the number of the bombarding particles hitting the target. The yield production of nuclei for any energy, E can be expressed as a function of the cross section as [8]:

$$Yield = I(\phi n) H (1 - e^{-\lambda t}) \int_{E_{out}}^{E_{in}} \sigma(E) \left(-\frac{dE}{dx}\right)^{-1}$$

Where: Y is the activity in (Bq) of the product nuclei. I: current of projectile in ( $\mu\text{A}$ ).  $\phi$  : the flux is  $\sim 10^{12}$  to  $10^{14} \text{ n/cm}^2 \cdot \text{s}$ . n: umber of atoms per unit volume (N/A).N: Avogadro's number. A: the mass number of the target in (amu). H: isotopic abundance (or enrichment) of the target.  $\lambda$  : The decay constant of the prod-

uct= $\left(\frac{0.693}{t_{1/2}}\right)$  in ( $h^{-1}$ ). t: time of irradiation in

(h).  $-\frac{dE}{dx}$  : the stopping power,  $\sigma(E)$  : Cross section at energy E in (mb).

**Results and Discussion**

Gallium radioisotopes  $^{66}\text{Ga}_{35}$  ( $t_{1/2} = 9.4\text{h}$ ),  $^{67}\text{Ga}_{36}$  ( $t_{1/2} = 78\text{h}$ ) and  $^{68}\text{Ga}_{37}$  ( $t_{1/2} = 68\text{min}$ ) are commonly used for diagnostic studies using (PET) and (SPET), it can be produced via various reactions induced by charged particle as shown in Table 1.

There are three routes for the accelerator production of  $^{66}\text{Ga}_{35}$  by proton projectile via  $^{66}\text{Zn}_{36}(p,n)^{66}\text{Ga}_{35}$ , and  $^{67}\text{Zn}_{37}(p,2n)^{66}\text{Ga}_{35}$  reactions as shown in Figures (1-a), and (2-a) respectively; also by deuteron bombardment via  $^{66}\text{Zn}_{36}(d,2n)^{66}\text{Ga}_{35}$  reaction as shown in (4-a).comparing the three reactions mentioned

above, the optimum reaction to produce  $^{66}\text{Ga}$  is  $^{66}\text{Zn}_{36}(p,n)^{66}\text{Ga}_{35}$  reaction with a production yield equals to  $7.4\text{MBq (mCi)/}\mu\text{Ah}$  (Figure 6), high cross section (700mb), at area of optimum

energy range  $E_p = (8 \rightarrow 14)\text{MeV}$ , and with no impurities. The obtained value is in a good agreement with the value obtained from ref. [2], with  $E_p = (6 \rightarrow 15)\text{MeV}$  because of the highly

isotopic abundance (27.9%) [9] of  $^{66}\text{Zn}$  in a natural Zinc matrix target. The yields of  $^{67}\text{Zn}_{37}(p,2n)^{66}\text{Ga}_{35}$  and  $^{66}\text{Zn}_{36}(d,2n)^{66}\text{Ga}_{35}$  reactions are  $0.093\text{MBq mCi/}\mu\text{Ah}$  and  $3.97\text{MBq(mCi)/}\mu\text{Ah}$  with optimum energy range  $E_p = (5 \rightarrow 15)\text{MeV}$  and  $E_p = (10 \rightarrow 15)\text{MeV}$  respectively.

The optimum production of  $^{68}\text{Ga}$  is via  $^{68}\text{Zn}_{38}(p,n)^{68}\text{Ga}_{37}$  reaction with a yield of  $30.34\text{MBq (mCi)/}\mu\text{Ah}$  (Figure 8), high cross section (1060mb), optimum energy range  $E_p = (8 \rightarrow 12)\text{MeV}$ , and with a minimum impurity 0.002% of  $^{67}\text{Ga}$ . This value is agree with value obtained from ref. [6] with

$E_p = (4 \rightarrow 13)MeV$  because of the highly isotopic abundance of  $^{68}Zn$  (27.9%) compared with  $^{67}Zn$  (4.1%) [9] in a natural Zinc matrix target. The results of the optimum energy range is less than 30MeV, hence the data obtained in the present work has a good agreement with some calculations made by researches. At which Gallium isotopes could conveniently yield at cyclotron energy for a medium *via* proton irradiation on natural Zinc target. However, calculations in the recent practical data of cross sections *via* deuteron irradiation on natural Zinc might be added a new production route for radio Gallium isotopes [4].

### Conclusions

The accelerators production of Gallium radioisotopes has suitable physical properties, since it have intermediate or short half-life, so it is widely used in medical applications in imaging techniques and radiotherapy. In the present work, the evaluation of cross sections for the reactions induced by proton and deuteron particles give a good agreement with experimental calculations taken from IAEA which done by different authors. The high enrichment target is used to reduce the radio impurities due to the activation of impurities, but this radio impurity does not disappear. These impurities can be removed only by using enriched target and/or by a careful selection of the effectively charged particle energy range in the target. From the results, we concluded the optimum reactions to product  $^{66}Ga$ ,  $^{67}Ga$  and  $^{68}Ga$  from  $^{66}Zn_{36}(p,n)^{66}Ga_{35}$ ,  $^{68}Zn_{38}(p,2n)^{67}Ga_{36}$ ,  $^{68}Zn_{38}(p,n)^{68}Ga_{37}$  and  $^{68}Zn_{38}(d,2n)^{68}Ga_{37}$  reactions respectively. Another reactions are not desirable to product  $^{66}Ga$ ,  $^{67}Ga$  and  $^{68}Ga$ , because of minimum product yield and the production of isotopic impurities appear in reaction process. We found from the results that Gallium-68 is the best isotope to be used in medical application because it have a short half-life and high cross sections.

**Acknowledgment:** We wish to express our deep thanks and gratitude to the International Atomic Energy Agency (IAEA) to allow access to the experimental nuclear data libraries and made available.

### References

- [1] International Atomic Energy Agency (IAEA), "Cyclotron Produced Radionuclides: Physical Characteristics and Production Methods", Technical Reports Series, No.468, P.2, Vienna (2009).
- [2] Rowshanfarzad P., Jalilian A. R., Sabet M., Akhlaghi M., "Production and quality control of  $^{66}Ga$  as a PET radioisotope", Iran, Journal of Radiation Research, Vo.2, No.3, pp: 149-158, (2004).
- [3] Amir R. J., Pejman R., Mahsheed S., Ali R. K., Amir A. R. , " Development of  $^{66}Ga$  oxine complex; a possible PET tracer", Nukleonika, Vol.51, No.3, pp: 155–159, (2006).
- [4] Khandaker M. U., Haba H. and Abu Kasim H., "Cyclotron produced  $^{67}Ga$ , a potential radionuclide for diagnostic and therapeutic applications", AIP Conference Proceedings 1659, 030009; doi: 10.1063/1.4916857, (2015).
- [5] Ochab E., Petelenz B., Szałkowski M., Ruszel M., Sroka A., Bogdali P., "Upgrade of the equipment for production of gallium isotopes from proton-activated zinc targets", Institute of Nuclear Physics Polish Academy of Sciences, (2014).
- [6] Mukesh K. P., John F. B., Huailei J., Alan B. P., Timothy R., " Cyclotron production of  $^{68}Ga$  via the  $^{68}Zn(p,n)^{68}Ga$  reaction in aqueous solution", American Journal of Nuclear Medicine and Molecular Imaging, Vol.4, No.4, pp.303-310, (2014).
- [7] <https://www-nds.iaea.org/exfor/endl.htm>
- [8] Qaim S. M., "Nuclear data for medical applications: an overview, Radiochim Acta. Journal. Vol.89, PP.189-196, (2001).
- [9] Krane K. S., "Introductory Nuclear Physics", John Wiley and Sons, PP.378-379, (1988).
- [10] Szelecsenyi F., Suzuki K., Kovacs Z., Takei M. and Okada K., "Investigation of the  $natZn(p,x)^{62}Zn$  nuclear process up to 70MeV: a new  $^{62}Zn/^{62}Cu$  generator", Journal of Applied Radiation and Isotopes, Vol.58, p.377, UK (2003).
- [11] Hermanne A., "Evaluated cross section and thick target yield data of Zn+P processes for practical applications", Journal of ARI, Vol.49, No. 1005, (1998).

- [12] Hermanne A., Walravens N. and Cicchelli O., "Optimization of isotope production by cross section Determination", conference on Nuclear Data for Sciences and technology, Juelich, p.616, Germany (1991).
- [13] Levkovskij V. N., "Activation cross section nuclides of average masses ( $A=40-100$ ) by protons and alpha-particles with average energies ( $E=10-50$  MeV)", Book: Levkovskij, Act. Cs. By Protons and Alphas, Moscow, (1991).
- [14] Tarkanyi F., Szelecsenyi F., Kovacs Z. and Sudar S., "Excitation Functions of Proton Induced Nuclear Reactions on Enriched  $^{66}\text{Zn}$ ,  $^{67}\text{Zn}$  and  $^{68}\text{Zn}$  Production of  $^{67}\text{Ga}$  and  $^{66}\text{Ga}$ ", Journal of Radiochimica Acta, Vol.50, p.19 Germany, (1990).
- [15] Hille M., Hille P., Uhl M. and Weisz W., "Excitation Functions of (p,n) and (a,n) Reactions on Ni, Cu and Zn", Journal of Nuclear Physics, Section A, Vol.198, p.625, Netherlands (1972).
- [16] Howe H. A., "(P, N) Cross Sections of Copper and Zinc.", Journal of Physical Review, Vol.109, p.2083, USA, (1958).
- [17] Skakun E. A., Utenkov S. N., Bondarenko V. N., Goncharov A. V., Mishchenko V. M., Sukhostavets V. I. and Shebeko K. V., "Cross sections and rates of the thermonuclear reactions  $^{64}\text{Zn}(p,g)^{65}\text{Ga}$  and  $^{66}\text{Zn}(p,g)^{67}\text{Ga}$ ", Journal of Bulletin of Russian Academy of Sciences - Physics, Vol.72, p.376, USA (2008).
- [18] Zelecsenyi F., Boothe T. E., Takacs S., Tarkanyi F., Tavanoe., "Evaluated cross section and thick target yield data bases of Zn+p processes for practical applications", Journal of Applied Radiation and Isotopes, Vol.49, p.1005, UK (1998).
- [19] Little F. E. and Lagunas-Solar M. C., "Cyclotron Production of  $^{67}\text{Ga}$ . Cross Sections and Thick-Target Yields for the  $^{67}\text{Zn}(p,n)$  and  $^{68}\text{Zn}(p,2n)$  Reactions", Journal of Radiochimica Acta, Vol.60, p.57, Germany (1993).
- [20] Hermanne A., "Evaluated cross section and thick target yield data of Zn+P processes for practical applications", Private communication Name. Hermanne (1997).
- [21] Krivonosov G. A., Ekhichev O. L., Nemashkalo B. A., Storizhko V. E. and Chirt V. K., "Radiative-Capture Cross Sections of Nuclides of Intermediate Atomic Weight for Low-Energy Protons", Journal of Izv. Rossiiskoi Akademii Nauk, Ser. Fiz., Vol.41, Issue.10, p.2196, Russia (1977).
- [22] Szelecsenyi F., Kovacs Z., Nagatsu K., Fukumura K., Suzuki K. and Mukai K., "Investigation of direct production of  $^{68}\text{Ga}$  with low energy multiparticle accelerator", Radiochimica Acta, Vol.100, p.5, (2012).
- [23] Mcgee T., Rao C. L., Saha G. B. and Yaffe L., "Nuclear Interactions of Sc-45 and Zn-68 with Protons of Medium Energy", Journal of Nuclear Physics, Section A, Vol.150, p.11, Netherlands (1970).
- [24] Hille M., Hille P., Uhl M. and Weisz W., "Excitation Functions of (P,N) and (A,N) Reactions On Ni, Cu and Zn", Journal of Nuclear Physics, Section A Vol.198, p.625, Netherlands (1972).
- [25] Steyn J. and Meyer B. R., "Production of  $^{67}\text{Ga}$  by deuteron bombardment of natural zinc", Journal of Applied Radiation and Isotopes, Vol.24, p.369, (1973).
- [26] Williams D. C. and Irvine J. W., "Nuclear excitation functions and thick-target yields: Zn+d and Ar-40(d, alpha)", Journal of Physical Review, Vol.130, No.1, p.265, USA (1963).
- [27] Vlasov N. A., Kalinin S. P., Ogloblin A. A., Pankratov V. M., Rudakov P., Serikov I. N. and Sidorov V. A., "Excitation functions for the reactions Mg-24(d,a)Na-22, Fe-54(d,a)Mn-52, Fe-54(d,n)Co-55, Zn-66(d,2n)Ga-66", Journal of Soviet Atomic Energy, Vol.2, p.189, USA (1957).
- [28] Nassiff S. J. and Munzel H., "Cross Sections for the Reactions  $^{66}\text{Zn}(D,N)^{67}\text{Ga}$ ,  $^{52}\text{Cr}(D,2N)^{52}\text{G}$  MN and  $^{186}\text{W}(D,2N)^{186}\text{RE}$ ", Journal of Radiochimica Acta, Vol.19, Issue.3, p.97, Germany (1973).
- [29] Gilly J. L., Henriet G. A., Preciosa A. M. and Capron P. C., "Absolute cross sections and excitation functions for (d,p) and (d,2n) reactions on  $^{55}\text{Mn}$ ,  $^{63}\text{Cu}$ ,  $^{65}\text{Cu}$ ,  $^{66}\text{Zn}$ , and  $^{68}\text{Zn}$  between 3 and 11.6 MEV", Journal of Physical Review Vol.131, No.4, p.1727, (1963).